



Marine Studies

The University of the South Pacific

Technical Report

A PRELIMINARY CHECKLIST OF THE MARINE
BENTHIC MACROALGAE OF ROTUMA

by

Antoine De Ramon N'Yeurt
Marine Studies Programme
University of the South Pacific
Suva
Fiji



PIM
GE
570
2
8891



Series
Number

93/2

ISSN 1018-2896

11/1

A PRELIMINARY CHECKLIST OF THE MARINE
BENTHIC MACROALGAE OF ROTUMA

by

Antoine De Ramon N'Yeurt
Marine Studies Programme
The University of the South Pacific
Suva, Fiji

August 1993

CONTENTS

Introduction	1
Methods	1
List of species	2
Discussion	10
Acknowledgements	11
References	11
TABLE I	13
TABLE II	13
FIGURE 1	12
FIGURE 2	14
FIGURE 3	14
FIGURE 4	14
FIGURE 5	14

INTRODUCTION

Rotuma (Fig. 1) is a small volcanic island about 465 km north of the main Fiji group, and is located at 12°30' South; 177°05' East. It occupies about 44 km², and lies in fair isolation from other island groups. Being six degrees North of Fiji, Rotuma experiences a rather more equatorial climate, and is quite distinct in terms of its structure, age and geography (Woodhall, 1987). It is encircled by a mostly fringing reef about 16 km² in area, which supports a varied algal flora.

To the author's knowledge, no phycological investigation of Rotuma took place prior to the present research, although the nearby Fijian algal flora has been fairly well studied (*vide* South and Kasahara, 1992). Hence this present checklist represents the first record of the Rotuman algal flora, which is dealt with in more detail by N'Yeurt (1993), South and N'Yeurt (1992) and South *et al.* (1993).

METHODS

Extensive algal collections were made on Rotuman reefs from January 1992 to May 1992, and also from December 1992 to February 1993, as part of the author's MSc Thesis project. Owing to the absence of diving, collections were limited to intertidal benthic habitats. Specimens were preserved in 5% formaldehyde in seawater, and shipped back to the University in Suva for eventual identification. Voucher specimens of all identified algae were deposited in the Phycological Herbarium, South Pacific Regional Herbarium, as either pressed specimens or slide collections. Statistical analysis on the results was carried out, including the computation of Jaccard Indices (Hoek, 1975) as a means of comparison between floras at different locations, namely between Fiji and Rotuma.

LIST OF SPECIES

A total of one hundred and six species are listed, comprising twenty-one Cyanophyceae, forty Rhodophyceae, nine Phaeophyceae and thirty-six Chlorophyceae. Taxa preceded by an asterisk (*) are new records for the Fijian algal flora, comprising forty-three taxa. A cross (+) indicates potentially new species.

The systematic order adopted, wherever possible, is that of Silva *et al.* (1987). Genera are listed alphabetically within each family, and species are listed alphabetically within each genus. The Cyanophyceae section broadly follows the classification in Bourrelly (1970) and most of the specimens were personally reviewed by B.A. Whitton during his visit to Fiji in March-April 1993.

SYSTEMATIC LIST CYANOPIHYCEAE CHROOCOCCALES Chroococcaceae

Aphanocapsa Nägeli
**Aphanocapsa* sp.
Gomphosphaeria Kützing
**Gomphosphaeria* sp.

Entophysalidaceae

Entophysalis Kützing
**Entophysalis* sp.

PLEUROCAPSALES Hyellaceae

Pleurocapsa Thuret ex Hauck
**Pleurocapsa* sp.
Dermocarpa P. Crouan et H. Crouan
**Dermocarpa* sp.
Oncobyrsa Meneghini
**Oncobyrsa* sp.

CHAMAESIPHONALES Chamaesiphonaceae

Chamaesiphon A. Braun et Grunow
**Chamaesiphon* sp.

Siphononemataceae

Siphononema Geitler

**Siphononema polonicum* Geitler

NOSTOCALES

Nostocaceae

Anabaena Bory de St. Vincent

**Anabaena* sp.

Oscillatoriaceae

Lyngbya C. Agardh

**Lyngbya aestuarii* (Mertens) Lieberman

Lyngbya "epiphytica" Hieron

Lyngbya majuscula (Dillwyn) Harvey

Oscillatoria Vaucher

**Oscillatoria* sp.

Schizothrix Kützing

**Schizothrix* sp.

Spirulina Turpin

**Spirulina tenerrima* Kützing

**Spirulina subtilissima* Kützing

Symploca Kützing

Symploca sp.?

Rivulariaceae

Calothrix C. Agardh

Calothrix sp.

Homoeothrix (Thuret) Kirchner

**Homoeothrix* sp.

Scytonemataceae

Tolypothrix Kützing

Tolypothrix sp.

Plectonema Thuret

Plectonema sp.

CHLOROPHYCEAE

ULVALES

Ulvaceae

Enteromorpha Link in Nees
Enteromorpha flexuosa (Wulfen) J. Agardh

CLADOPHORALES
Cladophoraceae

Cladophora Kützing
**Cladophora conferta* P. Crouan et H. Crouan
Rhizoclonium Kützing
**Rhizoclonium africanum* Kützing
**Rhizoclonium grande* Borgesen

SIPHONOCLADALES
Boodleaceae

Boodlea Murray et De Toni
Boodlea composita (Harvey) Brand
Struvea Sonder
Struvea anastomosans (Harvey) Piccone et Grunow ex Piccone

Siphonocladaceae

Boergesenia J. Feldmann
Boergesenia forbesii (Harvey) J. Feldmann
Cladophoropsis Borgesen
Cladophoropsis sundanensis Reinbold

Valoniaceae

Dictyosphaeria Decaisne ex Endlicher
Dictyosphaeria cavernosa (Forsskål) Borgesen
Valonia C. Agardh
Valonia aegagropila C. Agardh
Ventricaria Olsen and West
Ventricaria ventricosa (J. Agardh) Olsen et West

BRYOPSIDALES
Bryopsidaceae

Bryopsis Lamouroux
Bryopsis harveyana J. Agardh
**Bryopsis plumosa* (Hudson) C. Agardh

Caulerpaceae

Caulerpa Lamouroux

Caulerpa cupressoides (Vahl) C. Agardh

var. *lycopodium* Weber-van Bosse

*var. *lycopodium* (Weber-van Bosse) f. *elegans* (P. Crouan et H. Crouan) Weber-van Bosse

var. *mamillosa* (Montagne) Weber-van Bosse

Caulerpa racemosa (Forsskål) J. Agardh

var. *clavifera* (Turner) Weber-van Bosse

var. *peltata* (Lamouroux) Eubank

*var. *turbinata* (J. Agardh) Eubank

var. *uvifera* (Turner) J. Agardh

intermediate variety between vars. *turbinata* (J. Agardh) Eubank and *peltata* (Lamouroux) Eubank

Caulerpa serrulata (Forsskål) J. Agardh

*var. *boryana* (J. Agardh) Gilbert f. *occidentalis* (Weber-van Bosse) Yamada et Tanaka

Codiaceae

Codium Stackhouse

Codium arabicum Kützting

Codium bulbopilum Setchell

+**Codium* sp. nov.

Halimediaceae

Halimeda Lamouroux

Halimeda bikinensis W.R. Taylor

Halimeda cuneata Hering

Halimeda discoidea Decaisne

Halimeda macrophysa Askenasy

**Halimeda micronesica* Yamada

Halimeda opuntia (Linnaeus) Lamouroux

var. *opuntia* (Decaisne) J. Agardh

var. *hederacea* (Barton) Hillis

Halimeda simulans Howe

Halimeda taenicola W.R. Taylor

Halimeda tuna (Ellis et Solander) Lamouroux

Udoteaceae

Avrainvillea Decaisne

Avrainvillea nigricans Decaisne

+**Avrainvillea* sp. nov.

Chlorodesmis Harvey et Bailey

Chlorodesmis hildebrandtii A. Gepp et E.S. Gepp
**Chlorodesmis major* Zanardini
Rhipidosiphon Montagne
Rhipidosiphon javensis Montagne
Rhipiliopsis A. Gepp et E.S. Gepp
Rhipiliopsis sp.

DASYCLADALES
Polyphysaceae

Acetabularia Lamouroux
Acetabularia parvula Solms-Laubach

Dasycladaceae

Neomeris Lamouroux
Neomeris vanbosseae Howe

PHAEOPHYCEAE
ECTOCARPALES
Ectocarpaceae

Hincksia J.E. Gray
**Hincksia breviararticulata* (J. Agardh) P.C. Silva

SPHACELARIALES
Sphacelariaceae

Sphacelaria Lyngbye
Sphacelaria rigidula Kützing

DICTYOTALES
Dictyotaceae

Dictyopteris Lamouroux
Dictyopteris repens (Okamura) Borgesen
Dictyota Lamouroux
Dictyota friabilis Setchell
Dilophus J. Agardh
Dilophus radicans Okamura
Lobophora J. Agardh

Lobophora variegata (Lamouroux) Womersley

DICTYOSIPHONALES
Chnoosporaceae

Chnoospora J. Agardh

**Chnoospora minima* (Hering) Papenfuss

FUCALES
Sargassaceae

Sargassum C. Agardh

Sargassum polycystum C. Agardh

Turbinaria Lamouroux

Turbinaria ornata (Turner) J. Agardh

RHODOPHYCEAE
Bangiophycidae
BANGIALES
Erythropeltidaceae

Erythrotrichia J.E. Areschoug

Erythrotrichia carnea (Dillwyn) J. Agardh

Florideophycidae
ACROCHAETIALES
Acrochaetiaceae

Audouinella Bory

**Audouinella polyblasta* (Rosenvinge) J. Price, Lawson et John

BONNEMAISONIALES
Galaxauraceae

Actinotrichia Decaisne

Actinotrichia fragilis (Forsskål) Borgesen

GELIDIALES
Gelidiellaceae

Gelidiella Feldmann et Hamel
Gelidiella acerosa (Forsskål) Feldmann et Hamel
Gelidium Lamouroux
Gelidium pusillum (Stackhouse) Le Jolis

NEMALIALES
Liagoraceae

Liagora Lamouroux
**Liagora valida* Harvey

CORALLINALES
Corallinaceae

Cheilosporum (Decaisne) Zanardini
Cheilosporum spectabile Harvey ex Grunow
Fosliella Howe
Fosliella farinosa (Lamouroux) Howe
Jania Lamouroux
Jania adhaerens Lamouroux
**Jania rubens* (Linnaeus) Lamouroux
Mesophyllum Lemoine
**Mesophyllum mesomorphum* (Foslie) Adey
Porolithon (Foslie) Foslie
**Porolithon gardineri* (Foslie) Foslie

GIGARTINALES
Gracilariaceae

Gelidiopsis Schmitz
Gelidiopsis intricata (C. Agardh) Vickers
Gracilaria Greville
Gracilaria sp.

Solieriaceae

Meristotheca J. Agardh
**Meristotheca procumbens* P. Gabrielson et Kraft

RHODYMENIALES
Champiaceae

Champia Desvaux

Champia parvula (C. Agardh) Harvey

Rhodymeniaceae

Coelarthrum Børgesen

**Coelarthrum boergesenii* Weber-van Bosse

Coelothrix Børgesen

**Coelothrix irregularis* (Harvey) Børgesen

Rhodymenia Greville

**Rhodymenia divaricata* Dawson

CERAMIALES

Ceramiaceae

Centroceras Kützing

Centroceras apiculatum Yamada

Centroceras clavulatum (C. Agardh) Montagne

Ceramium Roth

**Ceramium codii* (Richards) Mazoyer

Ceramium mazatlanense Dawson

Ceramium vagans P.C. Silva

**Ceramium zacaе* Setchell et Gardner

Griffithsia C. Agardh

**Griffithsia subcylindrica* Okamura

Wrangelia C. Agardh

Wrangelia argus (Montagne) Montagne

Dasyaceae

Heterosiphonia Montagne

Heterosiphonia crispella (C. Agardh) Wynne

var. *laxa* (Børgesen) Wynne

Heterosiphonia subsecunda (Suhr) Falkenberg

Delesseriaceae

Hypoglossum Kützing

**Hypoglossum caloglossoides* Wynne et Kraft

Martensia Hering

Martensia elegans Hering

Rhodomelaceae

Amansia Lamouroux
Amansia glomerata C. Agardh
Bostrychia Montagne
Bostrychia tenella (Lamouroux) J. Agardh
Chondria C. Agardh
Chondria dasyphylla (Woodward) C. Agardh
Chondia minutula Weber-van Bosse
**Chondria sedifolia* Harvey

Herposiphonia Nägeli
Herposiphonia secunda (C. Agardh) Ambronn f. *tenella* (C. Agardh) Wynne
Laurencia Lamouroux
**Laurencia venusta* Yamada
Laurencia sp.
Polysiphonia Greville
Polysiphonia scopulorum Harvey
var. *scopulorum* (Harvey) Hollenberg

DISCUSSION

From the present list, it follows that forty-three fully identified taxa, or about 41% of the total Rotuman algal flora listed, represent new records for Fiji while two (*Avrainvillea* sp. nov., *Codium* sp.) represent new species, to be described elsewhere. Figure 3 shows the composition of the four main groups of Rotuman algae, while figure 4 compares the respective compositions of the Fijian and Rotuman algal floras. Figure 2 presents a summary of algal species at each station sampled; see N'Yeurt (1993) for further explanations. From Table I, It is immediately clear that the Fijian flora is much larger in terms of number of species; however the apparent impoverishment of the Rotuman flora is almost certainly a consequence of the lack of subtidal collections. Looking at Jaccard comparison indices (Table II; Figure 5), it appears that the Rotuman flora is quite dissimilar to the Fijian algal flora ($P_o = 19.9$); however once again this is largely a result of the vast size difference of the two floras and the absence of Rotuman subtidal collections, particularly Rhodophyta. It is pertinent to note that the two floras are most comparable in terms of Chlorophyta ($P_g = 26.9$), which are mostly found in intertidal habitats.

A particularly interesting observation is that one species of Rhodophyta found in Rotuma (*Meristotheca procumbens* P. Gabrielson et Kraft) represents the first observed record outside the type locality (Lord Howe Island; *vide* Gabrielson and Kraft, 1984), while a number of the Rotuman species are otherwise typically occurring in Northern Pacific waters. The latter biogeographic considerations are dealt with in more detail in N'Yeurt (1993).

The present preliminary research revealed quite interesting aspects of the Rotuman algal flora, providing an impetus for further work in this area. In particular, it would be most useful to obtain subtidal collections from Rotuman waters, especially Rhodophyta, in order to reassess

biogeographical comparisons of the Rotuman flora with neighbouring island groups.

ACKNOWLEDGEMENTS :

The author gratefully acknowledges financial support from the University of the South Pacific (URC grant # 0713-9106) and from the International Centre for Ocean Development (ICOD; Canada). I thank Dr. B.A. Whitton for his review of the Cyanophyceae, and Dr. G.T. Kraft for examining material of *Meristotheca*. I also wish to thank my supervisor, Professor G. Robin South, for unflinching support and encouragement throughout my research, and the staff of the Marine Studies Programme for technical assistance. Last but not least, heartfelt gratitude is expressed towards the kind people of Rotuma, without whose cooperation and support this study would not have been possible.

REFERENCES :

- Bourelly, P. 1970. *Les Algues D'eau Douce. Tome III. Les Algues Bleues et Rouges : Les Eugléniens, Peridiniens et Cryptomonadines*. Éditions N. Boubée & Cie, Paris. 512 pp.
- Gabrielson, P.W. and G.T. Kraft. 1984. The marine algae of Lord Howe Island (N.S.W.): the Family Solieriaceae (Gigartinales, Rhodophyta). *Brunonia* 7:217-251.
- Hoek, C. van den. 1975. Phytogeographic provinces along the coasts of the Northern Atlantic Ocean. *Phycologia* 14(4):317-330.
- N'Yeurt, A.D.R. 1993. *A Floristic Survey of the Intertidal and Shallow Benthic Marine Algae of Rotuma*. MSc Thesis, The University of the South Pacific [in preparation].
- Silva, P.C.; E.G. Meñez and R.L. Moe. 1987. Catalog of the benthic marine algae of the Philippines. *Smithson. Contrib. Mar. Sci.* 27:iv + 179pp.
- South, G.R. and H. Kasahara. 1992. A preliminary checklist of the benthic marine algae of the Fiji Islands, South Pacific. *Micronesica* 25(1):41-70.
- South, G.R. and A.D.R. N'Yeurt. 1993. Contributions to a catalogue of benthic marine algae of Fiji. II. *Caulerpa* and *Caulerpella* (Chlorophyta - Caulerpales). *Micronesica* [in review].
- South, G.R.; A.D.R. N'Yeurt and R.A. Raj-Prasad. 1993. Additions and amendments to the benthic marine algal flora of Fiji, including the island of Rotuma. *Micronesica* [in review].
- Woodhall, D. 1987. Geology of Rotuma. *Fiji Mineral Resources Dept. Bull.* 8:40pp+map.




Figure 1

LEGEND TO LOCATIONS ON MAP

- | | | | |
|-------------|------------------|--------------|------------|
| 1. Maka Bay | 2. 'Ahau | 3. Jölmea | 4. Mea |
| 5. Ropure | 6. Hoféa | 7. Lopta | 8. Oinafa |
| 9. Paptea | 10. Noa'tau | 11. 'Tua'koi | 12. Saolei |
| 13. Isilepi | 14. Hapmafau Bay | 15. Kelega | 16. Fapufa |
| 17. Losa | | | |

Table I. Numbers of Algal Species : Fiji and Rotuma

	Fiji *	Rotuma	New Records	Shared Species
All Species	326	106	43	66
Cyanophyceae	12	21	15	6
Chlorophyceae	99	36	11	28
Phaeophyceae	39	9	2	7
Rhodophyceae	176	40	15	25

* Excludes species occurring in Rotuma and not in Fiji.

Table II. Flora Comparisons : Fiji and Rotuma

	Jaccard Index (P)
All Species	19.9
Cyanophyceae	N/A
Chlorophyceae	26.9
Phaeophyceae	17.1
Rhodophyceae	13.3

Rotuman Algal Flora
 Composition At Each Station




Figure 2

Gr Bn Rd All

Rotuman Algal Flora
 Overall Composition




Figure 3

ALGAL TAXA NEW RECORDS

Rotuman and Fijian Algal Floras
 Overall Composition




Figure 4

Rotuman and Fijian Algal Floras
 Flora Comparisons : Jaccard Indices




Figure 5